

Recent Advances in the Field of Left-continuous T-norms

Sándor Jenei

Institute of Mathematics and Informatics
University of Pécs
Ifjúság u. 6, H-7624 Pécs, Hungary
jenei@ttk.pte.hu

Abstract

The recent advances in the research of left-continuous t-norms is summarized in this talk. The main focus is on construction methods, geometric description, and structural characterization. We point out further research directions and open problems.

Keywords: Left-continuous t-norm, Geometric description, Structural description, Characterization of subclasses.

1 Summary

The structure of continuous t-norms has been known since ages [19]. In many mathematical theories *left*-continuous t-norms have been used for a few decades without having a single (non-continuous) example at hand. After this long period – when no left-continuous t-norm¹ was known – the first example, the nilpotent minimum, was found by Fodor [2]. Even after finding this example many researchers believed that this is the only one, and a conjecture was published saying, roughly speaking, that the nilpotent minimum is the only left-continuous t-norm.

Next, a series of papers have appeared [11, 14, 15, 8, 7, 3, 1, 16, 5, 20, 6, 22], in which new construction methods provided the interested community with a huge number of new left-continuous t-norms. A state of the art is in [10].

On one hand, the complexity of different construction methods has reached a high level, therefore this line of research has become difficult to follow. On the other hand, the huge number of examples of left-continuous t-norms has called for a kind of systematization. One reply for that need is a comprehensive geometric characterization of residuated structures [9, 12], in par-

ticular, of left-continuous t-norms. This geometric description provides a kind of geometric view of algebraic phenomena, thus it makes the field of left-continuous t-norms, and its construction methods much more understandable. A recent result along this direction is [17], which resulted in another construction method in [18].

Concerning structural characterization, little is known as of today. As it will be explained, it seems to be hopeless to give classification for the whole class of left-continuous t-norms. However this does not exclude the possibility of characterizing certain subclasses of them. Two recent results concerns the cancellative, and the regular cases, respectively [4, 21].

These are the focuses of the talk.

Acknowledgement

Supported by the Hungarian-American Enterprise Scholarship Fund (HAESF).

References

- [1] R. Cignoli, F. Esteva, L. Godo, F. Montagna, On a class of left-continuous t-norms, *Fuzzy Sets and Systems* 131 (2002), 283-296.
- [2] J.C. Fodor, Contrapositive symmetry of fuzzy implications, *Fuzzy Sets and Systems* 69 (1995) 141–156.
- [3] P. Hájek, Observations on the monoidal t-norm logic, *Fuzzy Sets and Systems* 132 (2002), 107–112.
- [4] R. Horčík, Algebraic Properties of Fuzzy Logics, PhD Thesis, Czech Technical University, Prague 2004.
- [5] S. Jenei, A characterization theorem on the rotation construction for triangular norms, *Fuzzy Sets and Systems*, *Fuzzy Sets and Systems*, 136 (2003), 283–289.

¹From now under left-continuous t-norms we mean left-continuous but not continuous t-norms.

- [6] S. Jenei, F. Montagna, A general method for constructing left-continuous t-norms, *Fuzzy Sets and Systems*, 136 (2003), 263–282.
- [7] S. Jenei, A note on the ordinal sum theorem and its consequence for the construction of triangular norms, *Fuzzy Sets and Systems*, 126 (2002), 199–205.
- [8] S. Jenei, F. Montagna, A proof of standard completeness of Esteva and Godó's logic *MTL*, *Studia Logica*, 70 (2002), 184–192.
- [9] S. Jenei, Geometry of Associativity Theory and Application, *Proceedings of the 25th Linz Seminar on Fuzzy Set Theory (Fuzzy Logics and Related Structures)*, (Linz, Austria), 2005, 63–66.
- [10] S. Jenei, How to construct left-continuous triangular norms state of the art *Fuzzy Sets and Systems*, 143 (2004), 27–45.
- [11] S. Jenei, New family of triangular norms via contrapositive symmetrization of residuated implications, *Fuzzy Sets and Systems* **110** (1999), 157–174.
- [12] S. Jenei, On the geometry of associativity, *Semigroup Forum* (in press)
- [13] S. Jenei, On the structure of rotation-invariant semigroups, *Archive for Mathematical Logic*, 42 (2003), 489–514.
- [14] S. Jenei, Structure of left-continuous triangular norms with strong associated negations. (I) Rotation construction, *Journal of Applied Non-Classical Logics* 10 (2000), 83–92.
- [15] S. Jenei, Structure of left-continuous t-norms with strong associated negations. (II) Rotation-annihilation construction, *Journal of Applied Non-Classical Logics*, 11 (2001) 351–366.
- [16] S. Jenei, Structure of Girard monoids on $[0, 1]$, in: E.P. Klement, S. E. Rodabaugh, (Eds.), *Topological and Algebraic Structures in Fuzzy Sets*, Kluwer Academic Publishers, Dordrecht, (A Handbook of Recent Developments in the Mathematics of Fuzzy Sets, Trends in Logic, vol 20.) 2003, pp. 277–308.
- [17] K.C. Maes and B. De Baets, On the structure of left-continuous t-norms that have a continuous contour line, *Fuzzy Sets and Systems* 158 (2007), 843–860.
- [18] K.C. Maes and B. De Baets, The triple rotation method for constructing rotation-invariant t-norms, *Fuzzy Sets and Systems* (in press)
- [19] P.S. Mostert, A.L. Shields, *On the structure of semigroups on a compact manifold with boundary*. *Ann. Math.*, 65 (1957), 117–143.
- [20] D. Smutná, *On a peculiar t-norm*, *Busefal* 75 (1998), 60–67.
- [21] T. Vetterlein, Regular left-continuous t-norms (submitted)
- [22] Y. Ouyanga, J. Fanga, J. Lic, A conditionally cancellative left-continuous t-norm is not necessarily continuous, *Fuzzy Sets and Systems* 157 (2006), 2328–2332.